



# TritonGrip TE™ Nitrile Gloves

TritonGrip TE™ black, low-profile gloves feature exceptional tactile sensitivity. These textured nitrile gloves are suitable for all environments where barrier protection and user comfort is important. These gloves are 5 mil thick to reduce hand fatigue and enhance comfort. They are powder free, non-sterile and single use. No latex was used in the manufacturing of this product.

## SPECIFICATIONS

Material	Nitrile
Color	Black
Length	9.5 in
Cuff thickness	5 mil
Palm thickness	3 mil
Finger thickness	5 mil

## TENSILE STRENGTH

Before aging	45 MPa min
After aging	47 MPa min


## ELONGATION

Before aging	677%
After aging	626%

## Ordering Information

121-47133	Small, 100/bx
121-47134	Medium, 100/bx
121-47135	Large, 100/bx
121-47136	X-Large, 90/bx
121-47137	XX-Large, 90/bx

Meets NFPA 2018 Guidelines



## Call for your quote!

For more information on pricing and promotions, please call Customer Service at 800.558.6270 or visit [buyemp.com](http://buyemp.com)



# TritonGrip TE™ Nitrile Gloves

Tested for use with chemotherapy drugs using ASTM D6978.

Test Chemotherapy Drug and Concentration	Minimum Breakthrough Detection Time (Minutes)
Bleomycin Sulfate, 15 mg/ml (15,000 ppm)	No breakthrough up to 240 min.
Busulfan, 6 mg/ml (6,000 ppm)	No breakthrough up to 240 min.
Carboplatin, 10 mg/ml (10,000 ppm)	No breakthrough up to 240 min.
Carmustine (BCNU), 3.3 mg/ml (3,300 ppm)	WARNING: Not for use with Carmustine
Cisplatin, 1.0 mg/ml (1,000 ppm)	No breakthrough up to 240 min.
Cyclophosphamide (Cytoxan), 20 mg/ml (20,000 ppm)	No breakthrough up to 240 min.
Cytarabine, 100 mg/ml (100,000 ppm)	No breakthrough up to 240 min.
Dacarbazine (DTIC), 10.0 mg/ml (10,000 ppm)	No breakthrough up to 240 min.
Daunorubicin, 5 mg/ml (5,000 ppm)	No breakthrough up to 240 min.
Docetaxel, 10 mg/ml (10,000 ppm)	No breakthrough up to 240 min.
Doxorubicin Hydrochloride, 2.0 mg/ml (2,000 ppm)	No breakthrough up to 240 min.
Epirubicin (Ellence), 2 mg/ml (2,000 ppm)	No breakthrough up to 240 min.
Etoposide (Toposar), 20.0 mg/ml (20,000 ppm)	No breakthrough up to 240 min.
Fludarabine, 25 mg/ml (25,000 ppm)	No breakthrough up to 240 min.
Fluorouracil, 50.0 mg/ml (50,000 ppm)	No breakthrough up to 240 min.
Gemcitabine, 38 mg/ml (38,000 ppm)	No breakthrough up to 240 min.
Idarubicin, 1 mg/ml (1,000 ppm)	No breakthrough up to 240 min.
Ifosfamide, 50 mg/ml (50,000 ppm)	No breakthrough up to 240 min.
Irinotecan, 20 mg/ml (20,000 ppm)	No breakthrough up to 240 min.
Mechlorethamine, 1 mg/ml (1,000 ppm)	No breakthrough up to 240 min.
Melphalan, 5 mg/ml (5,000 ppm)	No breakthrough up to 240 min.
Methotrexate, 25 mg/ml (25,000 ppm)	No breakthrough up to 240 min.
Mitomycin C, 0.5 mg/ml (500 ppm)	No breakthrough up to 240 min.
Mitoxantrone, 2.0, mg/ml (2,000 ppm)	No breakthrough up to 240 min.
Paclitaxel (Taxol), 6.0 mg/ml (6,000 ppm)	No breakthrough up to 240 min.
Rituximab, 10 mg/ml (10,000 ppm)	No breakthrough up to 240 min.
Thiotepa, 10.0 mg/ml (10,000 ppm)	WARNING: Not for use with Thiotepa
Trisenox, 1 mg/ml (1,000 ppm)	No breakthrough up to 240 min.
Vincristine Sulfate, 1.0 mg/ml (1,000 ppm)	No breakthrough up to 240 min.
Fentanyl Citrate (100mcg/2mL)	No breakthrough up to 240 min.